Received: 26 March 2023 | Accepted: 17 May 2023

DOI: 10.1002/uro2.18

CONSENSUS

Indications, strategies, and development on prostate targeted biopsy: Report of the Panjiayuan Consensus Conference 2022

Gang Song[1] | Yajian Li[1] | Huimin Hou[2] | Yichen Wang[3] | Xuejuan Wang[4] | Shaoxi Niu[5] | Xiang Tu[6] | Hongliang Shen[7] | Zhien Zhou[8] | Yinbing Wang[9] | Ruiyi Yan[9] | Ning Xu[10] | Gejun Zhang[11] | Hailong Hao[12] | Shudong Zhang[13] | Ben Liu[14] | Bin Yang[15] | Baojun Wang[16] | Haifeng Wang[17] | Haifeng Huang[18] | Qian Zhang[19] | Baijun Dong[20] | Yonghong Li[21] | Jianbin Bi[11] | Weigang Yan[8] | Hongqian Guo[18] | Yuanjie Niu[22] | Guosheng Yang[17] | Ming Liu[2] | Qiang Wei[6] | Nianzeng Xing[1]

[1]Department of Urology, National Cancer Center/National Clinical Research Center for Cancer/Cancer Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China

[2]Department of Urology, Beijing Hospital, National Center of Gerontology, Institute of Geriatric Medicine, Chinese Academy of Medical Sciences, Beijing, China

[3]Department of Radiology, National Cancer Center/National Clinical Research Center for Cancer/Cancer Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China

[4]Department of Nuclear Medicine (PET-CT Center), National Cancer Center/National Clinical Research Center for Cancer/Cancer Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China

[5]Department of Urology, The Third Medical Center, Chinese PLA General Hospital, Beijing, China

[6]Department of Urology, Institute of Urology, West China Hospital, Sichuan University, Chengdu, China

[7]Department of Urology, Beijing Friendship Hospital, Capital Medical University, Beijing, China

[8]Department of Urology, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China

[9]Eight-year Medical Doctor Program, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China

[10]Department of Urology, Urology Research Institute, The First Affiliated Hospital, Fujian Medical University, Fuzhou, China

[11]Department of Urology, The First Hospital of China Medical University, Shenyang, China

[12]Department of Urology, Shanxi Cancer Hospital, Affiliated Cancer Hospital of Shanxi Medical University, Taiyuan, China

[13]Department of Urology, Peking University Third Hospital, Beijing, China

[14]Department of Urology, The First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou, China

[15]Department of Urology, Shanghai Tenth People's Hospital, Tongji University School of Medicine, Shanghai, China

[16]Department of Urology, Chinese PLA General Hospital/PLA Medical School, Beijing, China

[17]Department of Urology, Shanghai East Hospital, School of Medicine, Tongji University, Shanghai, China

[18]Department of Urology, Nanjing Drum Tower Hospital, The Affiliated Hospital of Nanjing University Medical School, Nanjing, China

[19]Department of Urology, Peking University First Hospital, Beijing, China

Gang Song, Yajian Li, Huimin Hou, Yichen Wang, Xuejuan Wang, Shaoxi Niu, Xiang Tu, Hongliang Shen, Zhien Zhou, Yinbing Wang, Ruiyi Yan, Ning Xu, and Gejun Zhang contributed equally to this study.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.
© 2023 The Authors. UroPrecision published by John Wiley & Sons Australia, Ltd on behalf of Higher Education Press.